

A Review on Groundwater Pollution

Hussein Ilaibi Zamil Al-Sudani

Petroleum Technology Department. University of Technology. Baghdad. Iraq

Abstract

Groundwater is one of the essential components in the development of any area. It is the primary potable, agricultural and industrial source of water. In the past, it was thought that layers of rocks and soils filter groundwater. The first signs of contamination of groundwater extracted from wells may be detected. The objective of this review is reviews researches of groundwater pollution and contamination as well as pollution types and the effects of groundwater contamination and pollution on public health. Generally, groundwater pollution studies include the scientific understanding of biological, chemical and physical processes which control contaminants fate and movement in the underground environment. High chemical concentration in drinking water can pose a health hazard. Epidemiological studies have shown that the low quality of drinking water as the main transport route has been responsible for many waterborne diseases. Several microorganisms and synthetic chemicals have the potential to pollute groundwater. Drinking water containing bacteria and viruses can lead to diseases such as hepatitis or cholera. The serious health effects of lead are known learning difficulties in children; problems in the nerves, kidneys, liver and pregnancy risks. Protection measures are more straightforward and less costly than corrective measures for groundwater contamination. The choice of appropriate therapeutic technique depends on site-specific factors and often takes into account clean-up goals based on human health and environmental protection.

Keywords: Groundwater, Contamination and Pollution, Review

I. INTRODUCTION

Water is essential for life and all economic activities. It is used for domestic, industrial and agricultural purposes. Having sufficient water in sufficient quantity and quality contributes to maintaining health. The availability of water of good quality is essential to prevent diseases and to improve the quality of life. The use of water increased due to an increase in human population and activities [1]. Groundwater is one of the essential components in the development of any area. It is the primary potable, agricultural and industrial source of water. In 2003, it was estimated that groundwater holds nearly 50% of the drinking water supply, 40% of the demand for industrial water, and 20% of the water used for irrigation [2]. Globally, more than a third of the water used by humans comes from groundwater. In rural areas, the ratio is higher: more than half of all drinking water worldwide is supplied from groundwater [3].

Groundwater pollution can also result from innumerable standard practices, such as the use of fertilizers and pesticides as well as disposal of human, animal and agricultural waste [4]. Globally, many researchers have conducted a study on the quality of groundwater and pollution sources affected by the industrial and natural process [5]. This concept has been developed from the vulnerability of groundwater. Therefore the most crucial part of the assessment of the risks of groundwater contamination [6, 7]. Quality changes of surface water, such as rivers or lakes, maybe due to contamination of groundwater. In the past two decades, awareness of groundwater pollution and contamination has been grown up [8]. Groundwater contamination effects become widely. Seven million people of the United States drink contaminated water yearly. The contaminated water caused by vomiting, rash, hepatitis, diarrhea closed all around the U.S. beaches. Groundwater pollution can lead to low drinking water quality, loss of water supply, degraded surface water systems, high cleaning costs, rising costs of alternative water supply and / or potential health problems [9]. The objective of this study is to review the researches of groundwater pollution and contamination as well as pollution types and the effects of groundwater contamination and pollution on public health. Some of the groundwater detection and remediation methods discusses in this review.

II. THEORY

Groundwater is the important potable, agricultural and industrial source of water. Therefore the ability to predict the conduct of chemical pollutants in the flowing groundwater is vital to a reliable assessment of hazardous or risks arising from groundwater pollution problems and the design of practical and effective mitigation techniques. Reliable prediction and quantification of pollutant movement can only be made if we understand the processes that control contaminant transport. These include: [10]

- 1- Physical, chemical and biological reactions that affect their soluble concentrations in groundwater,
- 2- Hydrodynamic dispersion
- 3- Advection.

The most groundwater contamination problems are: [4]

- 1- Prevent the introduction of contaminants to an aquifer
- 2- Predict movement of the contaminants if they are introduced as well as remove it to protect the biosphere effectively. Because of the importance of groundwater, difficulty and expense in addressing groundwater supplies, steps are often taken to prevent initial



contamination. These steps may include protection of the entire aquifer, as well as the area around the head of the surface well, from unintentional contamination [12, 13]. Studies of groundwater contamination generally include: [14, 15]

- The scientific understanding of biological, chemical and physical processes which control contaminants movement and fate in the subsurface environment;
- The mathematical representation in transport models to predict the movement of contaminants;
- The determination of different model parameters in the field and the laboratory using different methods;
- The development of transport models to predict contaminants movement incase they are introduced;
- The development of management models which controlling and/or preventing the introduction of the contaminant into an aquifer and determining methodology of safe disposal of hazardous wastes, and
- The development of methodology regarding contaminants removal to the extent necessary to protect the biosphere effectively.

Pollution of water sources occurs through point-to-point pollution and is not a source [16]. Pollution from an anonymous source is more ambiguous and can not be linked to a specific point, such as the application of pesticides and fertilizers in agricultural fields [17]. It has been shown that agricultural land, sewage, runoff and runoff from urban areas and animal waste are responsible for increasing the number of microorganisms found in water [16, 18, and 19]. The characters of water pollution point source are: a- Pollution can be accurately determined [20], and b- Limited real extent. A water pollution problem occurred when substances added to water to affect their chemical composition and caused threaten to human health and the ecosystem [21]. In addition to chemical contamination, groundwater quality can be further compromised by increasing levels of parameters that affect water quality, such as the concentration of hydrogen ions (pH), electrolysis (E.C.), turbidity, and microbiological content [22].

There are many routes by which groundwater are contaminated:

- Inter- aquifer leakage and irrigation returns [23].
- Leached from soluble solids at the surface and leaking from broken sewer pipes or lines [22].
- Percolation of liquid sprayed over land [24].
- Landfill leachate [25].
- Septic and sewerage effluent discharge [19].

Historically, groundwater has been a source of high quality and relatively clean, drinking water, needing little if any or no treatment [27].

III. TYPES OF GROUNDWATER POLLUTION

Groundwater is exposed to various sources of pollution due to industrial revolution together with the lack of appreciation of chemicals and their potential impact on land and water bodies.

The presence of pollution within groundwater is a significant challenge for delineation and identification. Leakage from chemical distribution infrastructure and petrochemicals, for example, pipelines and sewage collection systems such as sewage tanks, urban sewage channels and pipelines are a few examples of the real-life of unknown underground pollution. Products of mining activities and industrial complexes, which were stored in or underground without any rule to control the contamination of land, was one of the most challenging and difficult problems associated with pollution and management over the last hundred years [8].

The source pollutants characteristics which need to be identified include:

- The spatial locations of sources.
- The activity duration of sources which clarify the time of sources became active.
- The rate of injection of pollutant sources which determines the flow of contaminants from each source [8].

1.1 Groundwater Nitrate Contamination: Inorganic contamination of the most significant concern in groundwater is nitrate ions, which usually occur in aquifers near rural and suburban populations. Although uncontaminated groundwater generally has nitrate-nitrogen levels of less than 2 ppm, nitrate in groundwater originates mainly from four sources:

- Use of nitrogen fertilizers, inorganic and animal manure.
- Atmospheric deposition.
- Human sewage deposited in septic systems.
- Cultivation of the soil [33].

Approximately 12 million tones of nitrogen per year are used as fertilizer in agriculture within the United States, and the production of manure has contributed about 7 million tones or more. In most cases, nitrogen forms reduction in the soil is oxidized to nitrate, which then migrates to groundwater, where it dissolves in water, and it is diluted because nitrate removal from well water is costly. Water contaminated with high levels of nitrate is typically not used for human consumption, at least in public health [34].

1.2 Acidification: Acidified precipitation is a recognized and widespread phenomenon. The phenomenon of 'acid rain' has been known for over a century and is caused mostly by the release of oxides of sulphur and nitrogen into the atmosphere. Acid precipitation affects groundwater considered to be small. Other sources of acid in groundwater include an interaction of natural water-rock; contamination with industrial acids; and degradation of other contaminants [35].

1.3 Contaminated Land: Contaminated land presents the risk of groundwater contamination if the contaminants can migrate to the aquatic environment and continue to have potentially harmful concentrations. The consequences of pollution from human activities in the past and present have led and continue to lead to severe contamination of groundwater [36, 37, 38]

1.4 Heavy Metals: In the past, there has been a great interest in the fate of heavy metals due to the application of sewage sludge on agricultural land and from landfills. Additional sources of heavy metals include mining and smelting, burning of fossil fuels, metalworking, electronic and chemical industries, war and military training, and the launch of sports. In recent years, the recognition of potential toxic effects and long-term contaminants of heavy metal pollutants has focused on the protection of soil and groundwater [29].

1.5 Landfill: The potential landfills impact on groundwater quality hinges on the continuing debate between 'dilute and disperse' and 'containment' landfill design strategies. Understanding of landfill impact on groundwater requires knowledge of the composition of the landfill leachate. In general, leachate composition depends on the waste type, landfill design and practices, analytical procedures and timing. Most of the preceding references relate to the land filling of a mixed household which typically contains a significant organic fraction, is biodegradable and is subject to significant long term consolidation settlement due to low initial density and biodegradation effects. However, individual industrial solid wastes are typically landfilled in 'mono-disposal' facilities where only one or possibly two types of waste material are disposed of. Examples of these sites are deposits of minerals and mining industry wastes (e.g. colliery shale, coal tailings, quarry fines) and pulverized fuel ash (P.F.A.) from coal-fired power stations. Relatively few references were found on mono-disposal landfilling [39]. Pollution of dumpsite (an engineering landfill) identified as a significant threats of groundwater resources [40].

1.6 Microbiological Contaminants: Microbiological contamination of groundwater is derived from wastewater from humans or animals. The large variety of pathogens that may be present in sewage includes pathogenic bacteria, viruses and protozoa. These contaminants can represent a potentially severe threat to public health if they are present in a water supply. Microbiological contaminants may enter the subterranean environment by

leaking sewage, digging sinks, septic tanks, soaking liquids, mines used as a road to disposal, landfills or land-based drainage as fertilizer. The groundwater flow and degree of pollution subject primarily to the loading of liquid waste and groundwater exposure to surface-derived pollution. Previous studies on the movement of microbiological pollutants in the aquifer environment were more commonly associated with groundwater contamination incidents and to a lesser degree with investigative surveys, both of which concentrated on bacteriological and to a lesser degree on viral contamination [29]. Refuse (domestic and industrial solid waste) is defined as non-useful and undesirable material originating from human activities and not free-floating [29]. Poor management of solid materials has led to many catastrophic effects such as aesthetics, environmental hazards and pollution. Groundwater pollution may also occur due to the potential contamination of leachate from waste [41].

IV. REMEDIATION

Many organic and inorganic contaminants have been detected, mainly as a result of contamination from active and closed industrial facilities. It is anticipated that the legacy of past anthropogenic activities increasingly seen in groundwater, as today's contaminated land becomes tomorrow's groundwater pollution problem. Remediation is therefore rapidly becoming a necessary option, to restore or conserve groundwater quality and remove or contain contamination sources. The range of treatment required by the target concentration group is determined. There are two main categories of therapeutic objectives, the corrective target levels of the site and the non-specific target levels of non-site risks [29]. The presence of pollution in groundwater is a significant challenge for delimitation and identification. Treatment of the contaminated site requires an ideal decision-making system to accurately and efficiently identify the source of the pollutants [8].

1.1 Acidification: The possible environmental impacts of acid precipitation have been recognized in the last 25-30 years. A large number of researches have been published on the effects of acidification on streams and lakes, their environment, and on soil and more continuous. A recent review was done by Tickle, 1990. In contrast, relatively little work has been done on acidifications of groundwater, except for one central British Geological Survey review [94].

1.2 Contaminated Land: Following the identification of contaminated land sites, determination of the potential risks posed by the site is vital before proceeding with extensive site investigation or remedial activity. Risk assessment methodologies have received a great deal of attention in recent years, and a variety of approaches has been developed [38,43,44]. In general, future research on contaminated land is required to improve the understanding of the impact on groundwater quality, to classify risk and identify mitigation strategies. This

involves the collection of data on the extent of groundwater pollution from contaminated land, monitoring of the effectiveness of remediation procedures, as well as a better understanding of the processes controlling and influence the nature and extent of pollution [45].

1.3 Heavy Metals: The behaviour and mobility of trace metals in groundwater have been the subject of a growing number of research projects in the last decade. A detailed knowledge has been developed of the effects of pH and Eh conditions on the solubility, particularly to organic matter, clay minerals, hydrous iron and manganese oxides [46, 47].

1.4 Nitrate (NO_3^-): Nitrate (NO_3^-) groundwater contamination is a significant concern in all intensive agricultural areas (pollution from non-point sources). Reddy and Lin, (1999) [48] investigated the removal of nitrates from groundwater using a catalytic reduction process to selectively remove nitrate ions from groundwater, associated with the agricultural community. Palladium, platinum, and rhodium were used as three catalysts in this investigation. The catalytic reduction process described in this study is useful in removing NO_3^- from groundwater associated with non-source pollution. Puckett and Timothy, (2001) [49] investigated the transfer of nitrates in the aquifer concerning age, land-use practices and oxidation processes in central-western Minnesota. The researchers found that nitrate concentrations are high and behave like oxygen, decreasing with both increasing age and depth.

V. METHODS USED IN DETECTION OF GROUNDWATER POLLUTION

The increasing demand for groundwater needs to be monitored and regulated based on an overall assessment that includes the quality of groundwater, and not only the quantity, to ensure sustainable development and continuing use of the country's water resources [50,51]. Protection measures are more straightforward and less costly than corrective measures for groundwater pollution [52]. The precise and sufficient characterization of unknown sources of pollution in the groundwater system is the first critical step in the process of controlling and treating sub-surface pollution; it is more complicated related to pollution of surface water bodies which is relatively easier to detect. Some of the available methodologies for determining the source of groundwater pollution are reviewed here. However, it needs to be emphasized that often the efficiency of source identification depends on the availability and reliability of concentration field measurements and hydro-geologic information. So a control network designed to collect geochemical field measurements can help improve the efficiency of source identification.

The repetitive use of source identification models and the design monitoring network can be combined sequentially

to identify the source in an effective characterization methodology.

The characteristics of the pollutant source that need to be determined include:

- The sources in spatial locations.
- The activity duration of sources which clarify the time of sources became active.
- The rate of injection of pollutant sources that determines the flow of pollutants released from each source as a function of time [8].

Surface electrical resistivity was used as a tool to detect groundwater contamination. Water carries the pollutants through visible and invisible landscapes. At a local level, water-soluble components used in agriculture, industrial waste, dry waste sediments, may be discovered by water and produce groundwater pollution, which remains undetected until polluted water passes through a local well. The impact of pollution may appear in a downstream direction where river water is used for some vulnerable purposes such as irrigation [53].

There are four main assessment methods used to assess the risks of groundwater pollution with pesticides [28].

1. **Methods of indexing:** The methods of indexing are the most frequently used to assess groundwater pollution. Indexing methods use vital parameters that are near related to the entry of pollutants into groundwater, despite the vadose and saturated areas, for a comprehensive indicator of the characterization of pollution risks [28].
2. **Process-based methods:** Pollutants such as pesticides must pass through a series of physical, chemical and biological processes in the vadose region and the saturated zone. Simulation models use numeric models to describe related migrations and conversions. Under simulation models and to a certain extent, a numerical indicator can be calculated to characterize the risk of pollution [7].
3. **Statistical methods:** Based on observed pollution information, statistical methods can be used to detect the relationships between investigation data and pollution-related factors. Then, the evaluation indicators can be verified; each indicator with a value of relevance as analyzed by the statistical method is assigned [54,55].
- 4 - **Fuzzy comprehensive assessment method:** The fuzzy comprehensive assessment method requires first requires to make sure of the classification of pollution. Besides, the evaluation indicators should be selected, and then a standard value of each indicator under each contamination level must be collected. Second, the degree to which each sample belongs to each contamination level can be determined by merging the data into a relative membership score formula. In this step, the weight vector of the indices is also needed. Finally, by multiplying the

matrix of the relative optimal relative grade by the order matrix, a distinct value vector can be achieved, and the risk of pollution can be assessed according to this vector [56,57].

VI. WATER QUALITY AND PUBLIC HEALTH

Water quality and public health are linked in many ways [58]. Human life essentially depends on potable water [59]. Many theories have suggested that water has been responsible for the transmission of many waterborne diseases through bacterial contamination of drinking water, which is the most severe risk factor for the spread of diseases causing disease and death [60].

Despite global efforts and the availability of modern techniques for the production of safe drinking water, it has been reported that the transmission of waterborne disease is a matter of significant concern [61]. Pollution of drinking water during storage, lack of regulations and limited understanding and awareness among the population are documented [62,63].

The adverse effects of mechanical failure, human error or degradation of source water quality, even with the best treatment system and the disinfection process, can sometimes lead to water quality degradation [62,63,64]. Health risks can be caused by high chemical concentration in drinking water; however, in the case of bacteria contamination, there are only a small number of many that cause a danger to consumer health [65]. Epidemiological studies have shown that low quality of drinking water (referred to as feces, environmental pollution or chlorinated water) as the primary transmission route was responsible for many waterborne diseases [66].

Examples of such transport methods for waterborne diseases are:

1. Consumption of untreated, inadequately treated, or contaminated water directly or indirectly through food preparation.
2. Contaminated water contact with the body such as freshwater, seawater, swimming pool.

Several microorganisms and synthetic chemicals have the potential to pollute groundwater. Drinking water containing bacteria and viruses can lead to diseases such as hepatitis or cholera. Methemoglobinemia or Blue Baby Syndrome, a disease affecting infants, can be caused by drinking high nitrates water. Severe health effects of lead are known learning difficulties in children; neurological, kidney and liver problems and the risk of pregnancy [93].

VII. APPLICATIONS

Historically, groundwater contamination and pollution have been investigated all over the world using different methods and techniques. Some of these investigations and their results are shown below:

1. Jennings et al., (1998) [68] conducted a study on the toxicity of nitrates in groundwater. They showed that among infants less than one-year-old, pregnant women, people of all ages with reduced gastric acidity or hereditary lack of methemoglobin reeducates, those most vulnerable to exposure are those who use shallow wells for drinking water.
2. Hueb, (2001) [69] research groundwater contamination with chemicals such as cobalt. The study showed that (20%) have high levels of cobalt higher than 0.05mg.l^{-1} , after 25000 tests on wells implemented in Bangladesh, which is contrary to the (WHO, 1993) [22].
3. Water quality guidelines Merit and Press, (1997) [32] assessed water quality; high levels of calcium and magnesium were reported in aquifers in calcareous rocks.
4. Al-Sudani, (2003) [70] studied the Debagah basin in the north of Iraq where groundwater contamination was indicated with major anions, salinity and nitrate. Salinity and nitrate contamination was indicated in the southern and northern part of the basin.
5. Rosen et al., (2004) [71] investigated the prediction of groundwater nitrate contamination after the closure of the unlined sheep line in Carson, United States.
6. Kong et al., (2004) [72] surveyed the contamination of groundwater by pesticides in the potato growing area of Hebei and Shandong, and revealed the presence of aldicarb, phorate, and terbufos.
7. Lee et al. (2007) [73] discovered 13 types of organic chlorine pesticides in 56 shallow groundwater samples collected from an agricultural area in the Taihu Basin.
8. Huang et al., (2008) [74] found that organochlorine pesticides lightly pollute groundwater in the area of the Pearl River delta.
9. Akankpo and Igboekwe, (2011) [75] had monitored contamination of groundwater using surface electrical resistivity and geochemical methods. Five geo-electric soundings were implemented with two at the waste dump in Uyo, southwestern Nigeria. The results indicate that total (T.D.S.) and chloride values of water collected samples showing evidence of a quantitative contamination assessment in groundwater.
10. Al-Shamaa and Ali, (2011) [76] studied Badra-Jasan in Wasit province in Eastern Iraq. According to the surface and groundwater hydrochemical study in this basin according to salinity concentrations and major elements analyzed for 19 water samples, pollution of salinity and nitrate was identified. Pollution of both parameters has accrued in surface water and unconfined aquifer, while salinity pollution has been identified only in the confined aquifer of the Quaternary deposits and Muqdadiyah formation.

11. Lo Russo and Taddia, (2012) [77] studied a technical approach developed in Piemonte, northwest Italy, designed to protect drinking water wells. An essential aspect of this approach was to integrate the assessment of the vulnerability of the aquifer on a large scale while identifying the local WHPA delineation. This method has been successfully tested on a well for drinking water in the community at reasonable and adequate prices. However, for this method to be widely accepted, additional improvement is needed in some areas.

12. A semi-urban area of Croton Watershed (New York State, U.S.A.) provides 10% of New York City's drinking water. Unstable pollution in watersheds is a primary concern for managers because the water supply is currently unfiltered. The concentrations of sodium and dissolved chloride, while associated with the length of the road, stoichiometrically had more chloride than expected for pure road-salt dissolution. This displacement is likely to be due to the exchange of cations, and the absorption of sodium by wetlands in the watershed at Croton. The results showed that contamination in Hydrologic system of Croton must be addressed in current management policies and decision-making [78].

13. Identification of hydrogeochemical processes and Groundwater quality in Lagos University, Nigeria have been conducted by Odokoya, et al., (2013) [79]. The results of the study indicated that 50% of the water samples showed a pollution index higher than 1, and the lead contributed the largest percentage (37.8%) in the pollution index. This was followed closely by Mn, which contributed 29.3%. Concentrations of Al, Ni, Fe were 19.13%, 8.66%, 4.25% and 0.82% respectively.

14. Ehteshami et al., (2013) [80] analyzed the environmental impact of livestock facilities and nitrate leaching in groundwater in Ray- Iran. The Nitrate movement within soil and groundwater was simulated by LEACHN. By identifying different scenarios and analyzing performance sensitivity, they carefully studied the factors affecting groundwater pollution.

15. Al-Alwany, (2013) [81] studied the suitability of groundwater in Dabaa region – a western desert of Iraqi. Study results showed that toxic components of groundwater were effect on plants with mild to moderate effect of sodium concentration.

16. Coimbatore is one of the largest industrial and textile areas in southern India, where pollution of groundwater in an urban environment is a significant issue. Coimbatore groundwater system has deteriorated because of industrialization along with agricultural activities and rapid urbanization in surrounding areas [82].

17. Aliewi and Al-Khatib, (2015) [83] studied the Salfit District area, which is located in the northern part of the West Bank of Palestine. This area is located on recharge areas of karstified limestone aquifers. There are many pollutants in Salfit's aquifer recharge zone which

percolating and polluting into groundwater aquifers. The health of the Salfit population is directly threatened as results showed that water sources polluted using the Durov diagram. Microbiological analysis of potable water showed higher levels of total Coliforms.

18. Landfills are the most widely used method for municipal solid waste (M.S.W.) disposal method in China. However, these facilities have caused severe groundwater contamination due to the leakage of leachate. Han et al., (2016) [84] Studied and analyzed 32 scientific papers, a field survey and an environmental assessment report related to groundwater contamination caused by landfills in China. The groundwater quality in the vicinity of landfills was assessed as "very bad" by a total score (F.I.) of 7.85 by the grading method in China. Variety of pollutants consisting of 96 groundwater pollutants, 3 organic matter indicators, 2 visual pollutants and 6 aggregative pollutants had been detected in the various studies.

19. landfills method were used widely in China to dispose of municipal solid waste (M.S.W.). However, these facilities cause contamination of groundwater due to leakage of leachate. Han et al., (2016) [84] Studied and analyzed 32 scientific papers, field survey and environmental assessment report on groundwater pollution due to landfills in China. Groundwater quality in areas close to landfills was assessed as "very poor" according to (F.I.) of 7.85 through China grading method. A variety of contaminants consisting of 96 contaminated groundwater, 3 indicators of organic matter, 2 optical pollutants and 6 pollutants collected in different studies.

20. Majolagbe et al. (2017) [85] studied the potential vulnerability of groundwater in the surroundings to the pollution from Selous dumpsite (an engineering landfill) in Lagos, Nigeria was investigated. Physicochemical analyses revealed the groundwater with acidic pH which portend health risk, while microbiological examination showed the groundwater samples to exceed the permissible limits of World Health Organization guidelines total heterotrophic bacteria (<10 cfu /ml at 37°C) for drinking water. Various water quality indices were also used to show the variation of quality trends in the water samples collected.

21. Al-Sudani, (2017) [86] studied groundwater in Iraqi marshland area, where total dissolved salts of groundwater were very high concentration. The groundwater can't be used for human, agriculture and animal purposes due to the long distance of groundwater movements from recharge to discharge area beneath marshlands.

22. Al-Sudani, (2017) [87] studied Khan al-Baghdadi area which located in Anbar governorate in the west of Iraq. The results of groundwater salinity distribution as essential hydrochemical properties showed high concentration within carbonate aquifers extended in the area. The groundwater utilization can be used for

agriculture purposes although high concentrations of salinity.

23. Al-Sudani, (2018) [88] studied hydrogeological properties of groundwater in Karbala'a governorate in the middle west of Iraq, where several wells were polluted and contaminated with nitrate and significant as well as minor ions in the area of study.

24. Al-Sudani, (2019) [89] studied salinity pollution of groundwater in the south of Iraq. Depending on eighty-six wells, the aquifer formed by Dibdibba formation and alluvial fan deposits in the south of Iraq characterized by groundwater pollution with salinity while a small area located to the west of studied area was unpolluted zone.

25. Al-Sudani, (2019) [90] studied the groundwater system of Dibdibba sandstone aquifer in the south of Iraq, where multi wells were highly contaminated with several anions and cations as well as nitrate ion.

26. Al-Sudani, (2019) studied the hydrochemistry of groundwater in the northeast part of Anbar governorate in the west of Iraq.

VIII. CONCLUSIONS

The quality of groundwater is increasingly threatened due to agricultural, urban and industrial wastes that leak or are injected underlying aquifers. Groundwater is vulnerable to pollution as pollutants can enter from landfills and lakes used to store waste, chemical spills and underground storage tanks leakage, and hazardous waste sites improperly managed. Groundwater pollution can also result from innumerable standard practices, such as the use of fertilizers and pesticides; the disposal of human, animal and agricultural waste; and the use of chemicals. Groundwater pollution effects are extensive. Most severe effects of water pollution are human infectious disease in rural countries where sanitation facilities are seldom found. Protection measures are more straightforward and less costly than corrective measures for groundwater pollution. The choice of appropriate therapeutic technology depends on site-specific factors and often takes into account cleaning targets based on potential risks that protect human health and the environment. Many theories have suggested that water has been responsible for the transmission of many waterborne diseases through bacterial contamination of drinking water, which is the most severe risk factor for the spread of diseases. Diseases derived from chemical contamination of potable water are negligible compared to the number caused by microbial contamination. High chemical concentration in drinking water can lead to health risks.

IX. CONFLICTS OF INTEREST

The author declares that they have no conflicts of interest.

REFERENCES

- [1] H.I.Z. "Al-Sudani, *Hydrochemical Evaluation and Utilization of Groundwater*". Khanaqin Area, Diyala Governorate - East of Iraq. Iraqi Journal of Science, 59 (4C), 2018, 2279-2288.
- [2] S. S. D. "Foster and P. J. Chilton, *Groundwater: The Processes and Global Significance of Aquifer Degradation*," Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences, 358 (1440), 2003, 1957-1972. doi:10.1098/rstb.2003.1380
- [3] T. Harter, "Basic Concepts of Groundwater Hydrology. A.N.R. Publication 8083", FWQP Reference Sheet 11.1, University of California. U.S.A. 2015.
- [4] U.S. Environmental Protection Agency. Wellhead Protection: "A Guide for Small Communities, Chapter3": Ground Water Contamination, Office of Research and Development, Washington, DC 20460.144p. 1993.
- [5] S. Selvakumar, N. Chandrasekar and G. Kumar, "Hydrogeochemical characteristics and groundwater contamination in the rapid urban development areas of Coimbatore, India". Water Resources and Industry, 17, 2017, 26-33.
- [6] C.Z. Sun and S.S. Lin, "Review of ground water vulnerability concept and assessment", Jilin Geology, 19, 2000, 30-6
- [7] S. Foster, J. Chilton, M. Moench, F. Cardy and "M. Schiffler, *Groundwater in Rural Development: Facing the Challenges of Supply and Resource Sustainability*", World Bank Technical Paper 463, Washington DC. U.S.A. 2000.
- [8] M. Amirabdollahian and B. Datta, "Identification of Contaminant Source Characteristics and Monitoring Network Design in Groundwater Aquifers": An Overview. Journal of Environmental Protection, 4, 2013, 26-41.
- [9] "U.S. Environmental Protection Agency. Wellhead Protection: A Guide for Small Communities", Chapter 3: Ground Water Contamination, Office of Research and Development, Washington, DC 20460.144p. 1993.
- [10] "S. Foster, J. Chilton, M. Moench, F. Cardy and M. Schiffler, *Groundwater in Rural Development*": Facing the Challenges of Supply and Resource Sustainability, World Bank Technical Paper 463, Washington DC. U.S.A. 2000.
- [11] "S. Foster, R. Hirata", D. Gomes, M. D'Elia and M. Paris, Groundwater Quality Protection. World Bank, Washington DC, U.S.A., 2002, 103 p. doi:10.1596/0-8213-4951-1
- [12] A. A. Fadlelmawla and M. A. Dawoud, "An Approach for Delineating Drinking Water Wellhead Protection Areas at the Nile Delta", Egypt, Journal of Environmental Management, 79 (2), 2006, 140-149.
- [13] S. Y. Feng, S. Y. Huo, S. Z. Kang, Z. J. Tang and F. X. Wang, "Groundwater Simulation Using a Numerical Model under Different Water Resources Management Scenarios in an Arid Region of China", Environmental Earth Sciences, 62 (5), 2011, 961-971.
- [14] X. H. Wang, "Conceptual Design of a System for Selecting Appropriate Groundwater Models in Groundwater Protection Programs", Environmental Management, 21 (4), 1997, 607-615. doi:10.1007/s002679900053
- [15] W. Zhou, B. F. Beck, A. J. Pettit and B. J. Stephenson, "A Groundwater Tracing Investigation as an Aid of Locating Groundwater Monitoring Stations on the Mitchell Plain of Southern Indiana", Environmental Geology, 41 (7), 2002, 842-851.
- [16] G. Allenby and M. Margaret, "World, Environment and People. Science Press", N.S.W., Australia. 1993.
- [17] A.D. Viste and A.J. Rice, Major sources of groundwater contamination; "Assessing the extend of point and Non-point Contamination in Shallow. Aquifer System". Groundwater Contamination and Control. Edited by Uri Zoller. 1994.
- [18] C. Berka, D. McCallum and B. Wernick, "Resource Management and Environmental Studies". Department of Civil Engineering, University of Columbia, Vancouver, Bc, Canada. Paper presented at: Lower Fraser Basin in transition: Symposium and Workshop, Kwantlen College, Surrey, BC, Canada. 1995.
- [19] C.E. Boyd, Water Quality. "An Introduction. Kluwer Acad. Publisher", U.S.A. 330 p. 2000.

- [20] M.J. Hammer, "Water and Wastewater Technology", 3rd Edition, Published by Prentice-Hall Inc, Englewood Cliff, New Jersey. 1996.
- [21] A. J. Nathanson, "Basic Environmental Technology, Water Supply, Waste Management and Pollution Control". 2nd (eds) Published by Prentice-Hall Inc. 1997.
- [22] WHO: "World Health Organization. Guidelines for Drinking Water Quality", 2nd (eds), Vol.1. Recommendations. World Health; Geneva. 445p. 1993.
- [23] V.N. Reinhold, "Ground water Treatment Technology, 2nd Ed. Evan K Nyer". 1992.
- [24] M. J. Beckett, "Land Contamination", Contaminated Land, Problems and Solution. Edited by Tom Cairney. Published by Chapman and Hall. 1993.
- [25] E. G. Blight, "On Isolating a Landfill from the Surrounding Water regime", Microbiology of Landfill Sites, Eric Senor, 2nd ed, Published by C.R.C. Press. 1995.
- [26] U.S. Environmental Protection Agency. "Wellhead Protection: A Guide for Small Communities", Chapter 3: Ground Water Contamination, Office of Research and Development, Washington, DC 20460.144p. 1993.
- [27] "V.N. Reinhold, Ground water Treatment Technology", 2nd Ed. Evan K Nyer. 1992.
- [28] Y.Y. Zhao and Y.S. Pei, "Risk evaluation of groundwater pollution by pesticides in China: a short review". The 18th Biennial Conference of International Society for Ecological Modelling, Procedia Environmental Sciences, 13, 2012, 1739 - 1747.
- [29] A. Briffett, L. Houlden, S. Kirk, A. McMahon, P. Miles, R. Stephenson and E. Walker, "Groundwater Pollution Research Reviews. Geraghty and Miller International", Inc Conqueror House, Vision Park, Histon. Cambridge. 85 p. 1996.
- [30] C.A.J. Appelo and D. Postma, "Geochemistry, Ground water and Pollution", A.A. Balkama, Rotterdam. 536 p. 1999.
- [31] A.D. Eaton, L.S. Clesceri and A.E. Greenbury, "Standard methods for the examination of water and waste water", American Public Health Association, American Water Works Association and Water Environment Association, Washington D.C., U.S.A. 1995.
- [32] W. Merit and D. Press, "Water Quality and Pig Performance". Facts sheet; Ministry of Agriculture, Food and Rural Affairs, Ontario. 1997.
- [33] D.A.M. Barzinji and D.G.A. Ganjo, "Water Pollution", Limnological Investigations in Kurdistan region and Other part of Iraq. International Journal of Science, Environment and Technology, 3 (3), 2014, 776-799
- [34] C. Barid and M. Cann, "Environmental chemistry". 3rd Edition, W.H Freeman and Company press, pp 425. U.S.A. 2005.
- [35] M. Ford, J.H. Tellam and M. Hughes, "Pollution-Related Acidification in the Urban Aquifer, Birmingham", UK. Journal of Hydrology, 140, 1992, 297-312.
- [36] J.H. Tellam, "The Groundwater Chemistry of the Lower Mersey Basin Permo-Triassic Sandstone Aquifer System", U.K. - 1980 and Pre-Industrialisation Urbanisation. Journal of Hydrology, 161 (1-4), 1994, 287-325.
- [37] D.N. Lemer and J.H. Tellam, "The Protection of Urban Groundwater from Pollution. Journal of the Institute of Water and Environmental Management", 6 (1), 1992, 28-37.
- [38] M.P. Henton and P.J. Young, "Contaminated Land and Aquifer Protection. Journal of the Institute of Water and Environmental Management", 7(5), 1993, 539-547.
- [39] H. Robinson, J. Gronow, P.S. Durrant, M. Taylor, C.E. Reeve, P.G. Mackey, R. Mull and J.P.L. "Dearlove, Groundwater Protection in the U.K. - Assessment of the Landfill Leachate Source Term. Journal of the Institute of Water and Environmental Management", 6 (2), 1992, 229-236.
- [40] U.K. Singh, M. Kumar, R. Chauhan, P.K. Jha, A. Ramanathan and V. Subramanian, "Assessment of the impact of dumpsite on groundwater quality: a case study of the Pirana site in western". India. Environmental Monitoring Assessment 141, 2008, 309-21.
- [41] K. S. Makey, "Natural Buffers for Sludge Leachate Stabilization, Groundwater", 20 (4), 1982, 420- 429.
- [42] R.Muralidaran, G.Muthaiyan, M.Nicky wifred joseph, M.Sugumar, S.Suresh babu, "Analysis of Surface and Ground Water in Ranipet Industrial Area" SSRG International Journal of Civil Engineering 4.3 (2017): 54-57.
- [43] P.L. Walker, S. Munro, C.L. Hawkings, and F.E. Shepherd, "The Application of Risk Assessment to Contaminated Land - The British Gas Experience". Journal of the Institute of Water and Environmental Management, 8 (6), 1994, 607-614.
- [44] J.A. Cole, R.L. Norton and H.A.C. Montgomery, "Countering Acute Pollution Events – Procedures Currently being adopted in the United Kingdom" Water Science and Technology, 29 (3), 1994, 203-205
- [45] P. Bardos and H.J. "van Veen, Review of Longer Term or Extensive Treatment Technologies". Land Contamination and Reclamation, 4 (1), 1996, 19-36.
- [46] B.J. Alloway and A.P. Jackson, "The Behavior of Heavy Metals in Sewage Sludge Amended Soils". Science of the Total Environment, 1000, 1991, 223-257.
- [47] B.J. Alloway and D.C. Ayres, "Chemical Principles of Environmental Pollution, Blackie, Glasgow". 1994.
- [48] K.J. Readdy and J.M. Lin, "Nitrate removal from groundwater using catalytic reduction. Science direct research". 34 (3), 1999, 14.
- [49] L.J. Puckett and K.C. Timothy, "Transport and Fate of Nitrate in a Glacial Outwash Aquifer in Relation to Ground Water Age, Land Use Practices, and Redox Processes". United States Geological Survey USGS. American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America. Journal of Environmental Quality 31, 2001, 782-796.
- [50] R.C. Leyland, "Vulnerability Mapping in Karst Terrains, Exemplified in the Wider Cradle of Humankind World Heritage Site". MS.c thesis University of Pretoria, South Africa. 2008.
- [51] N. Goldscheider, "Karst groundwater vulnerability mapping: application of a new method in the Swabian Alb", Germany. Hydrogeology Journal, 13, 2005, 555-564.
- [52] A. Margane, "Management, Protection and Sustainable Use of Groundwater and Soil Resources in the Arab Region". Guideline for the Delineation of Groundwater Protection Zones. Vol. 5, ACSAD-BGR Technical Cooperation Project. 2003.
- [53] A.O. Akankpo and M.U. Igboekwe, "Monitoring Groundwater Contamination Using Surface Electrical Resistivity and Geochemical Methods". Journal of Water Resource and Protection, 3, 2011, 318-324
- [54] S.Y. Chen, G.T. Fu, H.C. Zhou and G.L. Wang, "Fuzzy analysis model and methodology for aquifer vulnerability evaluation". Journal of Hydraulic Engineering. 7, 2002, 23-30.
- [55] S.Y. Chen and G.L. "Wang Fuzzy optimization interaction model for assessing groundwater pollution potential and case study". Journal of Dalian University of Technology. 39, 1999, 811-5.
- [56] Z. Fang, C.L. Xiao, X.J. Liang and B. Feng "Fuzzy comprehensive evaluation of the groundwater vulnerability in songnen plain". Journal of Jilin University (Earth Science Edition), 3, 2007, 546-50.
- [57] L.J. Zhang, Z.Y. Gong and X.T. "Sun Fuzzy comprehensive judgment of groundwater environmental vulnerability". Natural Sciences Journal of Harbin Normal University, 17, 2001, 109-12.
- [58] H.G. Brutland, "World Water Day Thursday 22 March. Help Make the difference". 2001.
- [59] W.C. Michiels and D.L.E. "Moyson, Bacteriological Analysis. Handbook of Water Analysis"; edited by Leo M.L.Nollet. Published by Marcel Dekker Inc. 2000.
- [60] UNEPA/WHO: United Nation Environmental Program Agency/ "World Health Organization. A practical Guide to the Design and Implementation of fresh water quality studies and monitoring". Edited by Jamie Bartram and Richard Balance. Publish on behalf of UNEPA/WHO. 1996
- [61] M. Stevens, S. McCounell, P. Nadehaum, M. Chapman, S. Ananthakumar and J. McNeil, "Drinking Water Quality and Treatment Requirements": A risk based Approach. Water, 12-16. 1995,
- [62] W.R. Mackenzie, N.J. Hoxie, M.E. Proctor, M.S. Gradus, K.A. Blair, "A massive outbreak in Milwaukee of cryptosporidium infection transmitted through public water supply". New Engineering Journal Med. 331 (3), 1994, 161-167.
- [63] P.A. Roefer, J.T. Monscivitz and D.J. Rexing, "The Las Vegas cryptosporidiosis out break. JAWWA", 88, 1996, 95-106.

- [64] E. E. Geldreich, "Microbial Quality of Water Supply in Distribution System". Publisher C.R.C. Press Inc, Co-operate N.W. Boca, Roton, Florida 33-41. 1996.
- [65] WHO:World Health Organization, "Guideline for drinking water quality. Recommendations". 4th Ed. Geneva, World Health Organization. 520 p. 2011.
- [66] F.G. Lee and J.L. Anne, "Public Health Significance of Waterborne Pathogens in Domestic Water Supplies and Reclaimed Water". 1993.
- [67] R.Pakuththarivaalan, D.Ramraj, S.Shyam sunder, M.Shyalaja, N.S.Ashik "Quality Analysis of Ground Water in Cuddalore District Neyveli Block using Conventional Method", International Journal of Engineering Trends and Technology (IJETT), V46(3),162-168 April 2017.
- [68] D. Jennings, S. Paul and D.W. Diane, Nitrates in Drinking water. "Washington State Department of Health (WSDOH)". An electronic media retrieved from the website: <http://www.doh.wa.gov/ehp/dw>. 1997.
- [69] J.A. Hueb, "Improving Water, Planning and Management Worldwide". New World Water. Publication of World Water Council, France. 2001.
- [70] H.I.Z. Al-Sudani, "Hydrogeological System of Debagah Basin In North of Iraq", Ph.D. Thesis , University of Baghdad, College of Science. 153 p. 2003.
- [71] M.R. Rosen, R.R. Reeves, S. Green, B. Clothier and N. Ironside, "Prediction of Ground water nitrate contamination after closure of an unlined sheep feedlot. Vadose Zone Journal". 3 (6), 2004, 990-1006.
- [72] D.Y. Kong, Z.L. Zhu, L.L. Shi, Z.J. "Shan and D.J. Cai, Effect of pesticides on groundwater under sweet-potato-based cropping systems in northern China". Journal Agro-Environ Sci, 23, 2004,1017-20.
- [73] B.H. Li, Z.Y. Ren, H.H. Chen, X.J. Cao and F. Liu, "Residues of organochlorine pesticides in shallow groundwater of agricultural region in Taihu basin", Journal of Agro-Environ Sci.26, 2007, 1714-8.
- [74] G.X. Huang, J.C. Sun, S. Wang, H.Y. Du, Y.D. Lu, B.F. Zhi, "Elementary research of organochlorine pesticide in groundwater of Pearl River delta". Journal Agro-Environ Sci, 27, 2008, 1471-5.
- [75] A.O. Akankpo and M.U. Igboekwe, "Monitoring Groundwater Contamination Using Surface Electrical Resistivity and Geochemical Methods". Journal of Water Resource and Protection, 3, 2011, 318-324
- [76] I.M. Al-Shamaa and B. M. Ali, "Hydrochemical Pollution of Groundwater in Badra-Jassan basin / Eastern of Iraq. Diyala Agricultural Sciences Journal". 3 (2), 2011, 669-679.
- [77] S. Lo Russo and G. Taddia, "Aquifer Vulnerability Assessment and Wellhead Protection Areas to Prevent Groundwater Contamination in Agricultural Areas": An Integrated Approach. Journal of Water Resource and Protection, 4, 2012, 674-685.
- [78] J.M. McKenzie, D.I. Siegel, L. K. Lautz, M. H. Otz, J. Hassett and I. Otz, "Water Quality, Contamination, and Wetlands in the Croton Watershed, New York", U.S.A. Open Journal of Modern Hydrology, 2, 2012, 7-14.
- [79] M. A. Odukoya, A.F. Folorunso, E.A. Ayolabi and E.A. Adeniran, Groundwater Quality and Identification of "Hydrogeochemical Processes within University of Lagos, Nigeria". Journal of Water Resource and Protection, 5, 2013, 930-940.
- [80] M. Ehteshami, A.S. Langeroudi and S. Tavassoli, "Simulation of Nitrate Contamination in Groundwater Caused by Livestock Industry (Case Study: Rey)". Journal of Environmental Protection, 4, 2013, 91-97.
- [81] A.A.M. Al-Alwany, "Suitability of groundwater of Dabaa region" – Iraqi western desert for agricultural use. Iraqi journal of civil engineering, 10 (1), 2013, 24-32.
- [82] V.S. Uday, A. Amar, P.S. Kunwar, D. Ratnakar and P.S. Netra, "Groundwater quality appraisal and its hydrochemical characterization in Ghaziabad (a region of indogangetic plain)", Utta Pradesh, India, Applied Water Science. 4, 2014, 145–157.
- [83] A. Aliewi and A.I. Al-Khatib, "Hazard and risk assessment of pollution on the groundwater resources and residents". health of Salfit District, Palestine. Journal of Hydrology: Regional Studies, 1(5), 2015, 10.
- [84] Z. Han, H. Ma, G. Shi, L. He, L. Wei and Q. Shi, "A review of groundwater contamination near municipal solid waste landfill sites in China", Science of the Total Environment 569 (570), 2016, 1255-1264.
- [85] A.O. Majolagbe, A.A. Adeyi, O. Osibanjo, A.O. Adams and O.O. Ojuri, "Pollution vulnerability and health risk assessment of groundwater around an engineering Landfill in Lagos", Nigeria. Chemistry International, 3(1), 2017, 58-68.
- [86] H.I.Z. Al-Sudani, "Groundwater Investigation in Iraqi Marshland Area". Diyala Journal for Pure Sciences. 13 (3), 2017, 12-29.
- [87] H.I.Z. Al-Sudani, "Hydrogeological study of Khan Al-Baghdadi area in Anbar governorate - West of Iraq. Diyala Journal for Pure Sciences". 13 (2), 2017, 192-207.
- [88] H.I.Z. Al-Sudani, "Hydrogeological Properties of Groundwater in Karbala'a Governorate – Iraq". Journal of University of Babylon, Engineering Sciences, 26(4), 2018, 70-84.
- [89] H.I.Z. Al-Sudani, "Salinity Pollution of Groundwater in South of Iraq". Journal of University of Babylon, Engineering Sciences, 27 (3), 2019, 13-24
- [90] H.I.Z. Al-Sudani, "Groundwater System of Dibdibba Sandstone Aquifer in South of Iraq". Applied Water Science. 9(4), 2019, 1-11.
- [91] H.I.Z. Al-Sudani, " Hydrochemistry of Groundwater in Northeast Part of Anbar Governorate – West of Iraq". Baghdad Science Journal. 16 (1), 2019, 88-96.
- [92] Desai Birva and Desai Hemangi, " Evaluation of Water Quality Index for Ground Water of Residential Area of Surat City, Gujarat, India" SSRG International Journal of Agriculture & Environmental Science 5.3 (2018): 1-5.
- [93] M.J. Pawari and S. Gawande, "Ground Water Pollution and Its Consequences". International Journal of Engineering Research and General Science, 3 (4), 2015, 773-776.
- [94] A. Tickle, "Acid Waters in the United Kingdom: Evidence, Effects and Trends". In Acid Deposition - Origins. Impacts and Abatement Strategies. J. W. Longhurst (ed). Springer Verlag. 1990.